

Oxysterols and retinal degeneration

Zhang, Xun; Alhasani, Reem Hasaballah; Zhou, Xinzhi; Reilly, James; Zeng, Zhihong; Strang, Niall; Shu, Xinhua

Published in:
British Journal of Pharmacology

DOI:
[10.1111/bph.15391](https://doi.org/10.1111/bph.15391)

Publication date:
2021

Document Version
Author accepted manuscript

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):

Zhang, X, Alhasani, RH, Zhou, X, Reilly, J, Zeng, Z, Strang, N & Shu, X 2021, 'Oxysterols and retinal degeneration', *British Journal of Pharmacology*, vol. 178, no. 16, pp. 3205-3219.
<https://doi.org/10.1111/bph.15391>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

Oxysterols and retinal degeneration

Xun Zhang¹, Reem Hasaballah Alhasani^{1,5}, Xinzhi Zhou¹, James Reilly¹, Zhihong Zeng⁴, Niall Strang²,
Xinhua Shu^{1,2,3*}

1 Department of Biological and Biomedical Sciences, Glasgow Caledonian University, Glasgow G4
0BA

2 Department of Vision Science, Glasgow Caledonian University, Glasgow G4 0BA

3 School of Basic Medical Sciences, Shaoyang University, Shaoyang, Hunan 422000, P. R. China

4 College of Biological and Environmental Engineering, Changsha University, Changsha, Hunan
410022, P. R. China

5 Department of Biology, Faculty of Applied Science, Umm Al-Qura University, Makkah, Saudi Arabia

* Corresponding authors: Xinhua.Shu@gcu.ac.uk

ABSTRACT

Retinal degeneration, characterized by the progressive death of retinal neurons, is the most common cause of visual impairment. Oxysterols are the cholesterol derivatives produced via enzymatic and/or free radical oxidation that regulate cholesterol homeostasis in the retina. Preclinical and clinical studies have suggested a connection between oxysterols and retinal degeneration. Here, we summarize early and recent work related to retina oxysterol-producing enzymes and the distribution of oxysterols in the retina. We examine the impact of loss of oxysterol-producing enzymes on retinal pathology and explore the molecular mechanisms associated with the toxic or protective roles of individual oxysterols in different types of retinal degeneration. We conclude that increased efforts to better understand the oxysterol-associated pathophysiology will help in the development of effective retinal degeneration therapies.

KEYWORDS

Retinal degeneration, cholesterol, oxysterol, toxicity, therapy

Abbreviations: 4 β -HC, 4 β -hydroxycholesterol; 7 α ,24S-diHC , 7 α ,24(S)-dihydroxycholesterol; 7 α ,25-diHC, 7 α ,25-dihydroxycholesterol; 7 α ,27-diHC, 7 α ,27-dihydroxycholesterol; 7-DHC, 7-dehydrocholesterol; 7-KC, 7-ketocholesterol; 7 α -HC, 7 α -hydroxycholesterol; 7 α / β -HC , 7 α / β -hydroxycholesterol; 7 β -HC, 7 β -hydroxycholesterol; 22R-HC, 22(R)-hydroxycholesterol; 20R,22R-diHC, 20R,22R-dihydroxycholesterol; 24S-HC, 24(S)-hydroxycholesterol; 24S,25-EC, 24(S),25-Epoxycholesterol; 25-HC, 25-hydroxycholesterol; 27-COOH, 3 β -hydroxy-5-cholestenoic acid; 27-HC, 27-hydroxycholesterol; ABCA1, ATP Binding Cassette Subfamily A Member 1; ABCG1, ATP Binding Cassette Subfamily G Member 1; ABCG4, ATP Binding Cassette Subfamily G Member 4; AMD, age-related macular degeneration; APOE, Apolipoprotein E; CYP, cytochrome P450; DHCR7, 7-dehydrocholesterol reductase; ER, endoplasmic reticulum; GCL, ganglion cell layer; INL, inner nuclear layer; INSIG, insulin induced gene; IPL, inner plexiform layer; LDLR, low-density lipoprotein receptor; LXR α and β , liver X receptor α and β ; NFL, nerve fibre layer; ONL, outer nuclear layer; oxLDL, oxidized LDL; RPE, retinal pigment epithelial; SCAP: Sterol regulatory element binding proteins cleavage-activating protein; SLOS, Smith-Lemli-Opitz syndrome; SREBP, sterol regulatory element-binding protein.

1 INTRODUCTION

Oxysterols are a group of oxygenated products from cholesterol or its precursors (Guillemot-Legris et al., 2016). A single oxysterol can be produced by an enzyme-mediated reaction, a non-enzymatic route, or from a combination of both processes (Figure 1). A sterol ring of cholesterol is oxidized by non-enzymatic routes including free radical and non-radical modification (Murphy and Johnson, 2008) and endogenous oxysterols, such as 7-ketocholesterol (7-KC, abbreviations and nomenclature listed in Table S1) and 7 β -hydroxycholesterol (7 β -HC), are also predominantly produced via non-enzymatic routes. A side chain of cholesterol is oxidized through enzymatic routes, typically catalysed by cytochrome P450 (CYP) enzymes. The CYP family members, including [CYP3A4](#), [CYP3A5](#), [CYP7A1](#), [CYP7B1](#), [CYP11A1](#), [CYP27A1](#), [CYP39A1](#) and [CYP46A1](#), are well characterised monooxygenases in oxysterol metabolism. CYP11A1 and CYP27A1 are localized to mitochondria, while the other CYP members are localized to endoplasmic reticulum (ER). CYP3A4 converts cholesterol to 4 β -Hydroxycholesterol (4 β -HC) and 25-Hydroxycholesterol (25-HC), while CYP3A5 converts cholesterol only to 4 β -HC (Bodin et al., 2002); CYP7A1 catabolizes cholesterol to 7 α -Hydroxycholesterol (7 α -HC) and catabolizes 7-Dehydrocholesterol (7-DHC) to 7-ketocholesterol (7-kC), while CYP7B1 oxidises 25-HC and 27-Hydroxycholesterol (27-HC) to, respectively, 7 α ,25-Dihydroxycholesterol (7 α ,25-diHC) and 7 α ,27-Dihydroxycholesterol (7 α ,27-diHC); CYP11A1 metabolizes cholesterol to 22(R)-Hydroxycholesterol (22R-HC) and from there to 20R,22R-Dihydroxycholesterol (20R,22R-diHC) then to pregnenolone; CYP27A1 converts cholesterol to 25-HC or to 27-HC (predominant); CYP46A1 oxidises cholesterol to 24(S)-Hydroxycholesterol (24S-HC), which can be metabolised to 7 α ,24(S)-Dihydroxycholesterol (7 α ,24S-diHC) by CYP39A1. Some CYP enzymes can also oxidize cholesterol precursors to form oxysterols, such as CYP46A1 that metabolises cholesterol precursor, desmosterol, to 24(S),25-Epoxycholesterol (24S,25-EC). Enzymatically and non-enzymatically formed oxysterols can be transported to the liver where they are further converted to bile acids, which facilitates the elimination of excess cholesterol from extrahepatic tissues. Oxysterols play an important role in a diverse range of cellular functions and

are involved in cancer, metabolic disorders and neurodegenerative diseases (Griffiths and Wang, 2019; Guillemot-Legaris et al., 2016).

Retinal degeneration occurs in a group of retinopathies that are characterised by the progressive death of retinal neurons. There are different types of retinal degeneration, mainly classified either as inherited or complex retinal disorders. Inherited retinal degeneration, such as retinitis pigmentosa, is caused by genetic defects, primarily affecting photoreceptor cells; whereas complex retinal degeneration involves a combination of genetic and environmental risk factors and includes conditions such as diabetic retinopathy, age-related macular degeneration and glaucoma. Cholesterol is enriched in the neural retina and in retinal pigment epithelial (RPE) cells and plays a critical role in maintaining retinal function (Pikuleva and Curcio, 2014). Major oxysterol-producing enzymes have been reported to be expressed in the retina and various oxysterols have been shown to be distributed in the retina (Fliesler and Bretillon, 2010; Pikuleva and Curcio, 2014). In this review we will discuss oxysterol metabolism in the retina and its function in retinal degeneration.

2 RETINAL STRUCTURE AND FUNCTION

The human retina is a 0.5mm-thick circular disc with a diameter of about 30-40mm that is positioned at the back of the eye. The retina is a highly organised light-sensitive structure which consists of a complex network of sensory neurons and associated cells which are tightly packed together (Figure 2A). The photoreceptor cells are the initial light detecting cells and their capability to detect light is dependent on their interaction with the RPE cells.

As shown in Figure 2A the RPE in the vertebrate eye lies between the outer segments of the photoreceptors and the choroidal blood supply. It is composed of a single layer of tightly-packed columnar hexagonal epithelial cells containing pigment granules (melanin) and other organelles involved in the digestion of photoreceptor outer segment membranes into phagosomes (Hu et al., 2008; Strauss, 2011). The melanosomes containing the pigment melanin are involved in the absorption of the light energy that strikes the retina. The RPE is aligned along Bruch's membrane through its basolateral membrane. Bruch's membrane provides a matrix essential for the interaction

of the RPE with blood flow in the capillaries of the choroid. The RPE therefore forms a tight-junction epithelium between the blood flow of the choroid and the outer photoreceptor segments, forming the outer blood-retinal barrier that enables selective permeability between the choroid and neurosensory retina (Steinberg, 1985; Strauss, 2011). This barrier is vital for the highly selective bi-directional processes involving the transport of nutrients from the blood to the photoreceptors (choroid to retina) and for the movement of water and metabolites from the retinal side to the blood (retina to choroid) (Strauss, 2011). The pigments in the RPE absorb light and reduce its scattering, thereby improving image resolution and helping to protect the retina from any large influx of light. The RPE cells act as storage reservoirs of metabolites and vitamins and are critical for maintenance of retinal adhesion (Katz and Gao, 1995).

The retina is composed of at least six types of neural cells (rods, cones, bipolar, horizontal, amacrine and ganglion cells) and one type of glial cell (Muller glia) derived from the neural ectoderm. These cells are compactly arranged in several layers including the outer nuclear layer (ONL), outer plexiform layer (OPL), inner nuclear layer (INL), inner plexiform layer (IPL), ganglion cell layer (GCL) and nerve fibre layer (NFL) (Figure 2 A) (Purves et al., 2001). The human retina comprises over 110 million rods and over 5 million cones (Remington, 2012) and these two types of photoreceptors are arranged in parallel fashion in the photoreceptor layer of the retina. Cones are responsible for photopic (bright illumination) central vision while the rods are responsible for scotopic (dim illumination) peripheral vision. Although rods outnumber cones by ~20 fold, cones are crucial for good visual acuity and colour discrimination. Human rods are highly sensitive, possessing the capacity to detect single photons, whereas each cone requires receipt of more than 100 photons to generate a comparable response (Mustafi et al., 2009). Phototransduction, in which a photon of light is changed to an electrical signal, takes place in the outer segments of the rod and cone photoreceptors. The electrical signals created by these retinal cells are then transferred via bipolar cells to the ganglion cells that convey the information to the brain through the optic nerve, resulting in the experience of vision (Remington, 2012). There is a striking difference in the circuitry

associated with rod and cone photoreceptors, specifically in the mode of transmission of signals from rods and cones to ganglion cells. Typically, a single cone cell communicates with a single bipolar cell that in turn transmits information to a single ganglion cell, whereas several rod cells can communicate with one bipolar cell, with several of the latter then feeding into a single ganglion cell. This convergence of signals by rods helps them to pool small signals in dim light, while the one-to-one relationship between cones-bipolar-ganglion cells maximizes visual acuity (Solomon and Lennie, 2007; Mustafi et al., 2009).

3 CHOLESTEROL HOMEOSTASIS IN THE RETINA

The retina has a well-regulated cholesterol input and output that maintains cholesterol homeostasis (Fliesler and Bretillon, 2010). The retina can directly take up cholesterol from the systemic circulation predominantly through a low-density lipoprotein receptor (LDLR)-mediated process in the RPE, which has LDLR localized to the basolateral side, facing the choroid (Tserentsoodolet et al., 2006). Local cholesterol synthesis also contributes to cholesterol input in the retina. Lin et al. (2016) demonstrated that *in situ* biosynthesis of cholesterol accounted for 72% of the cholesterol input in mouse retina. Excess cholesterol in the retina is either removed via the reverse cholesterol transport (RCT) and/or metabolized to oxysterols by CYP enzymes (Fliesler and Bretillon, 2010); however, quantitative contributions of the RCT and oxysterol formation to the removal of excess cholesterol in the retina remain elusive. Cholesterol efflux is the first step of the RCT pathway, which is mediated by acceptors (HDL and apolipoproteins) and ABC transporters (ATP Binding Cassette Subfamily A Member 1 ([ABCA1](#)), ATP Binding Cassette Subfamily G Member 1 ([ABCG1](#)) and [ABCG4](#)). Recent work from our laboratory and elsewhere has demonstrated effective cholesterol efflux in RPE and choroidal endothelial cells (Biswas et al., 2017; Lyssenko et al., 2018; Storti et al., 2017).

Proteins associated with regulation of cholesterol homeostasis (sterol regulatory element-binding protein, SREBP; Sterol regulatory element binding proteins cleavage-activating protein, SCAP, insulin induced gene, INSIG; liver X receptor α and β , [LXR \$\alpha\$](#) and [LXR \$\beta\$](#)) are expressed in the retina (Zheng et al., 2012). SREBP-2 is the major regulator of cholesterol homeostasis. When intracellular

cholesterol is at a low level, SCAP escorts SREBP-2 from endoplasmic reticulum (ER) to the Golgi apparatus, where SREBP-2 is cleaved by two proteases, resulting in the release of the amino-terminal of transcription factor functional domain and upregulation of cholesterol synthesis and uptake associated genes; when cholesterol level is high, cholesterol binds to SCAP and triggers SCAP physically interacting with INSIG, leading to retaining SCAP in the ER and blocking SREBP-2 associated transcriptional activation (DeBose-Boyd et al., 1999; Hua et al., 1993; Yang et al., 2002). Furthermore, oxysterols inhibit cholesterol synthesis by binding to INSIG and preventing SREBP-2 processing (Adams et al., 2004; Radhakrishnan et al., 2007). Additionally, oxysterols can regulate cholesterol homeostasis by activating the LXR signal pathway and enhancing cholesterol efflux (Fu et al., 2001; Janowski et al., 1999; Lehmann et al., 1997).

4 OXYTEROLS IN THE RETINA

4.1 Expression of major oxysterol-producing enzymes in the retina

Some oxysterol-producing enzymes, such as CYP family members, have been shown to express in the retina (Figure 2B). Guarneri et al (1994) first reported that CYP11A1 was localized to ganglion cells and cells of the inner nuclear layer of rat retina; CYP11A was also shown to be similarly localized in hamster retina, although here its expression in the cells of the inner nuclear layer was higher than that in rat retina (Jaliffa et al., 2005). Lee et al (2005) examined expression of CYP27A1 in monkey retina and found that CYP27A1 was strongly expressed in photoreceptor inner segments and was also expressed at low levels in the RPE, choroid, ganglion cells, nerve fibres and Müller cells.

CYP46A1 had higher mRNA levels in the neural retina and low levels in the RPE in bovine samples (Bretillon et al., 2007). CYP46A1 was strongly localized to ganglion cells and was also present in the inner nuclear layer of rat retina (Bretillon et al., 2007; Ishikawa et al., 2016); CYP46A1 also had similar localization in mouse retina (Ramirez et al., 2008). While CYP27A1 and CYP46A1 were also expressed in the human RPE cell line, ARPE-19 (Biswas et al., 2017) and monkey endothelial cell line, RF/6A (Biswas et al., 2018). In the human retina, mRNA of *CYP11A*, *CYP27A1* and *CYP46A1* has been detected in the neural retina and RPE. CYP11A was localized to ganglion cells and the cells of the

inner nuclear layer in both human and rodent retinas. CYP11A1 was also expressed in Bruch's membrane, the RPE and photoreceptor inner segments in human retina. CYP27A1 was predominantly localized to Bruch's membrane, RPE, photoreceptor inner segments, ganglion cells and nerve fibre layer. CYP46A1 had a localization in human retina similar to that of CYP27A1, although was expressed at a lower level (Zheng et al., 2012). Early work in the same lab demonstrated that CYP27A1 was expressed at a higher level in human neural retina (403-510 fmol/mg protein) than was CYP46A1 (58-63 fmol/mg protein), while CYP27A1 was also expressed in human RPE at 1110-2060 fmol/mg protein, about 2.7-4.0-fold higher than in the neural retina (Heo et al., 2011; Liao et al., 2011).

4.2 Distribution of oxysterol in the retina

Based on the expression of oxysterol-producing genes (discussed above) and high level of oxidative stress (Wright et al., 2010), oxysterols in the retina are generated via enzymatic and radical reactions in a way similar to that seen in other tissues. A number of studies have examined oxysterols in the retinas of different species and found that oxysterols are present in the RPE, photoreceptor inner segments, ganglion cells and nerve fibre layer (Moreira et al., 2009; Rodriguez et al., 2014; Rodriguez and Fliesler, 2009) (Figure 2B). Moreira et al. (2009) examined oxysterols in primate retinas using high-pressure liquid chromatography-mass spectrometry (HPLC-MS) and immunohistochemistry. 7-KC, which is predominantly generated by non-enzymatical oxidation (Figure 1 and ref Anderson et al., 2020), was present in the neural retina at 1-1.5 pmol/nmol free cholesterol and in RPE/choroid at 5-8 pmol/nmol free cholesterol. Immunostaining showed that 7-KC was localized to monkey choriocapillaries, RPE and Bruch's membrane (Moreira et al., 2009). Rodriguez and Fliesler (2009) investigated the effect of photodamage on generation of the oxysterols 7-KC and 7 α / β -hydroxycholesterol (7 α / β -HC) in rat neural retina. They reported that light-treated rats had an approximately 6-fold increase in 7-KC (control rats: 1.6 pmol/nmol; treated rats: 12.7 pmol/nmol cholesterol) and a 50-fold increase in 7 α / β -HC (control rats: trace only; treated rats: 25.0 pmol/nmol cholesterol), indicating increased production; further immunohistochemistry showed that the

increased 7-KC was localized to the RPE, photoreceptor inner segments and ganglion cells of photodamaged retinas (Rodriguez and Fliesler, 2009). In aged mice, a significantly high level of 7-KC was accumulated beneath the RPE (Indaram et al., 2015). 7-KC was also examined in bovine retina and was found to be lower in the neural retina (25 pmol/mg protein) than in the RPE (41 pmol/mg protein); a further two 7-KC-derived oxysterols, 3β ,27-dihydroxy-5-cholesten-7-one (7-KCh-27OH) and 3-hydroxy-5-cholesten-7-one-26-oic acid (7-KCh-27COOH), were detected at low levels in bovine RPE (7-KCh-27OH: 1 pmol/mg protein; 7-KCh-27COOH: not quantified) but not in the neural retina (Heo et al., 2011). Measurement of 7-KC in monkey neural retina and RPE/choroid demonstrated that 7-KC level was increased during aging and was 3-4 times higher in the RPE/choroid (1-68 pmol/nmol cholesterol) than in the neural retina (1-18 pmol/nmol cholesterol). Similarly, 7-KC in aged human RPE (2483.8 pmol/nmol cholesterol) was around 150 times higher than that in the neural retina (16.5 pmol/nmol cholesterol) (Rodriguez et al., 2014). Drusen, cholesterol- and lipid-rich extracellular deposits underneath the RPE, also contain an extremely high level of 7-KC (734.9 pmol/nmol cholesterol) (Curcio et al., 2005; Rodriguez et al., 2014). However, 7-KC in human plasma is unaffected by aging and is not associated with age-related macular degeneration (AMD), the leading cause of blindness in aged population (Lin et al., 2018), suggesting increased *in situ* synthesis of 7-KC in neural retina and RPE with aging.

Using gas chromatography-mass spectrometry (GC-MS), Mast et al. (2011) examined the CYP enzymes-mediated oxysterol production: 24(S)-Hydroxycholesterol (24S-HC), produced by CYP46A1; 27-Hydroxycholesterol (27-HC) and 3β -hydroxy-5-cholestenoic acid (27-COOH) (produced by CYP27A1); 22(R)-Hydroxycholesterol (22R-HC) and pregnenolone (produced by CYP11A1) in bovine and human retinas; they found abundant cholesterol (total) in the bovine neural retina (158 nmol/mg protein) and RPE (136 nmol/mg protein) but detected very low levels of oxysterols, including 24S-HC (total) at 0.036 nmol/mg protein in retina and at 0.021 nmol/mg protein in the RPE, 22R-HC (total) at 0.007 nmol/mg protein in neural retina and at 0.010 nmol/mg protein in RPE, and pregnenolone (total) at 0.019 nmol/mg protein in the neural retina and at 0.029 nmol/mg protein in

the RPE. However, 27-HC was undetectable in both the bovine neural retina and RPE, 27-HC oxidized 27-COOH (free) was present in both neural retina (0.025 nmol/mg protein) and RPE (0.09 nmol/mg protein), suggesting that when 27-HC is generated by CYP27A1, it might be immediately oxidised to 27-COOH by the same enzyme. In the bovine choroid, only 24S-HC (total) and 22R-HC (total) were detected at, respectively, 0.018 nmol/mg protein and 0.001 nmol/mg protein. Furthermore, significantly higher amounts of 27-COOH (free) was found in human neural retina (0.037-0.125 nmol/mg protein) than in the RPE (0.002-0.01 nmol/mg protein), while free 24S-HC and pregnenolone were detected at lower levels (0.001-0.004 nmol/mg protein for both) in the neural retina but was not detected at all in the RPE; free 22R-HC and 27-HC also were undetectable in both the neural retina and RPE (Mast et al., 2011; Liao et al., 2011).

5 LOSS OF OXYSTEROL-PRODUCING ENZYMES IS ASSOCIATED RETINAL PATHOLOGY

5.1 CYP27A1 and CYP46A1

Mutations in the *CYP27A1* gene cause cerebrotendinous xanthomatosis (CTX), which is a rare autosomal recessive metabolic disease that has systemic and neurological symptoms (Cruysberg et al., 1995; Dotti et al., 2001) and involves a variety of ocular abnormalities, including cataract, cholesterol-like crystals in the vitreous humour, premature retinal senescence, retinal vessel sclerosis, optic disk paleness, drusen and RPE defects (Cruysberg et al., 1995; Dotti et al., 2001; Morgan et al., 1989). *Cyp27a1*^{-/-} mice have been extensively characterized. Such mice do not exhibit the classic CTX symptoms such as cataract, xanthomas in tendon and brain, and atherosclerosis (Honda et al., 2001; Repa et al., 2000; Rosen et al., 1998). However, retinal pathology in *Cyp27a1*^{-/-} mice was first characterised in Dr Pikuleva's laboratory (Omarova et al., 2012). *Cyp27a1*^{-/-} mice demonstrated defects in retinal cholesterol homeostasis and developed cholesterol deposits beneath the RPE, neovascularization and Müller cell activation (Omarova et al., 2012).

Cyp46a1^{-/-} mice demonstrated a greater than 40% reduction in cholesterol turnover in the brain but had a similar level of hepatic cholesterol and bile acid synthesis (Lund et al., 2003). Saadane et al (2019) examined the systemic and retinal cholesterol metabolism and retinal phenotypes. The

systemic cholesterol metabolism of the *Cyp46a1*^{-/-} mice did not change significantly but did exhibit dysregulation of retinal cholesterol homeostasis. Vision function testing found no differences in either scotopic (rod function) or photopic (cone function) electroretinogram (ERG) between wildtype and *Cyp46a1* deleted mice; however, *Cyp46a1*^{-/-} mice demonstrated venous beading, tortuosity, increased vascular permeability and microglial activation in the retina (Saadane et al., 2019). Since CYP27A1 and CYP46A1 are the main enzymes for cholesterol metabolism in the retina, the same group also examined retinal pathology of *Cyp27a1*^{-/-}/*Cyp46a1*^{-/-} mice and found that these animals had significantly higher levels of retinal cholesterol compared to *Cyp27a1*^{-/-} or *Cyp46a1*^{-/-} alone; the mutant animals also developed abnormal retinal vasculature and had macrophage/microglial activation and increased oxidative stress in the retina. ERG tests showed that both scotopic and photopic ERGs were significantly lower in male double knockout mice when compared to the wildtype male mice (Saadane et al., 2014). Sterol-O-acyltransferase 1 (SOAT1) is responsible for cholesterol esterification in the retina (Saadane et al., 2016), leading to production of cholesterol esters, which are a significant component of drusen (Curcio et al., 2005). Apolipoprotein E (APOE), a major apolipoprotein in the retina, regulates cholesterol homeostasis as an acceptor for cholesterol efflux or a ligand for LDLR (Mazzone and Reardon, 1994; Zhao and Mazzone, 1999). To further investigate the associated function of CYP27A1 and CYP46A1 with SOAT1 or APOE in cholesterol metabolism, Petrov et al. (2019) generated *Cyp27a1*^{-/-}/*Cyp46a1*^{-/-}/*Soat1*^{-/-} and *Cyp27a1*^{-/-}/*Cyp46a1*^{-/-}/*ApoE*^{-/-} mice, whose retinal structure and vasculature were normal. *Cyp27a1*^{-/-}/*Cyp46a1*^{-/-}/*Soat1*^{-/-} mice were found to have a normal retinal cholesterol level similar to that of *Soat1*^{-/-} mice, but the *Cyp27a1*^{-/-}/*Cyp46a1*^{-/-} mice had a high level of retinal cholesterol, suggesting a possible compensatory response in cholesterol synthesis. However, *Cyp27a1*^{-/-}/*Cyp46a1*^{-/-}/*ApoE*^{-/-} mice had a level of retinal cholesterol similar to that of *ApoE*^{-/-} mice. All *Cyp27a1*^{-/-}/*Cyp46a1*^{-/-}/*Soat1*^{-/-} mice exhibited markedly decreased scotopic ERGs, with decreased photopic ERGs seen only in the male mice. *Soat1*^{-/-} mice had similar trends of decreased ERGs to that of *Cyp27a1*^{-/-}/*Cyp46a1*^{-/-}/*Soat1*^{-/-} mice, indicating that cholesterol esterification is required for

retinal function. Only male *Cyp27a1*^{-/-}/*Cyp46a1*^{-/-}/*Soat1*^{-/-} mice had a significant decrease in both scotopic and photopic ERGs (Petrov et al., 2019).

5.2 DHCR7

7-dehydrocholesterol reductase (DHCR7) is the final enzyme of the Kandustsch -Russel pathway in the cholesterol biosynthesis pathway and converts 7-dehydrocholesterol (7-DHC) to cholesterol. Mutations of the *DHCR7* gene cause Smith-Lemli-Opitz syndrome (SLOS), which is an autosomal recessive metabolic disorder that is characterised by increased 7-DHC and reduced cholesterol in all tissues, malformation in multiple organs and mental retardation (Nowaceyk and Irons, 2012). SLOS patients also present with a wide range of ocular abnormalities, such as sclerocornea, cataract, paleness of optic discs and optic atrophy (Elias et al., 2003; Garry et al., 2010; Kretzer et al., 1981). Pathological analysis of a one-month-old SLOS male patient's eye samples showed generally normal retinal structure, suggesting that loss of DHCR7 may not affect early retinal development; nevertheless, there was extensive death of ganglion cells, disintegration of mitochondria in the RPE, and accumulation of cytoplasmic material underneath photoreceptors (Kretzer et al., 1981). SLOS children have been shown to have defects in rod and rod-derived functions, showing slow kinetics of phototransduction activation and deactivation, possibly due to dysfunction of membrane-associated phototransduction proteins (Elias et al., 2003). A follow-up study examined visual function by ERG testing of the same patient cohort with long-term supplementation with dietary cholesterol. Compared to the controls, patients exhibited a significant decrease in the rod response but no difference in the cone response, suggesting that additional dietary cholesterol may protect cone function in SLOS patients (Garry et al., 2010).

Retinal function of DHCR7 has been studied in rodent models. *Dhcr7* null mice have significantly increased 7DHC and markedly decreased cholesterol in serum and tissues. These mice also recapitulate some SLOS phenotypes, such as retardation of intra-uterine growth and craniofacial abnormality (Fitzky et al., 2001; Wassif et al., 2001). However, *Dhcr7* mutant mice only survive 18-24 hours after birth, limiting the application of the model for further phenotypic characterization. An

alternative SLOS rat model has been developed by treating normal rats with trans-N,N-bis[2-Chlorophenylmethyl]-1,4-cyclohexanedimethanamine dihydrochloride (AY9944), a selective inhibitor for DHCR7. AY9944-treated rats recapitulate SLOS phenotypes (Kolf-Clauw et al., 1996). When pregnant rats were fed with AY9944 from gestational day 6 till postnatal day (P) 28, and their offspring also received AY9944 treatment from birth to P28, the pups at P28 exhibited abnormal accumulation of 7-DHC and markedly reduced cholesterol in serum, liver, retina and brain when compared to control animals; however, there was no defect in retinal development or change in ERG function, indicating that loss of DHCR7 may not affect retinal development (Fliesler et al., 1999). When AY9944 treatment of the progeny was extended from 4 to 10 weeks, the offspring demonstrated retinal degeneration with shorter outer segments, reduced thickness of outer nuclear layer and decreased ERG amplitudes, along with abnormal ultrastructure of RPE cells, when compared to the control animals (Fliesler et al., 2004).

Retinal degeneration in AY994-treated rodents is supposed to be caused by 7-DHC-derived oxysterols (Fliesler and Xu, 2018). 7-DHC is a strong reactive lipid molecule, sensitive to free radical oxidation, so generation of oxysterols from 7-DHC is predominantly via nonenzymatic oxidation but also via enzymatic oxidation (Fliesler and Xu, 2018; Xu et al., 2008). Xu et al. (2010) firstly identified fourteen novel oxysterol species from 7-DHC via free radical oxidation and found the oxysterol mixture is toxic to Neuro2a cells at $\geq 10 \mu\text{M}$ possibly via inhibition of proliferation and induction of differentiation (Korade et al., 2010; Xu et al., 2010). 7-DHC derived oxysterols have been identified and are accumulated in blood, liver, brain and retina of AY9944-exposed rats, particular 7-KC in the AY9944-treated retina is more than 30 times higher compared to that of the controls (Xue et al., 2011, 2012). Intravitreal injection of 7-KC caused pan-retinal degeneration in normal rats, indicating the toxicity of 7-KC (Xu et al., 2012). Further *in vitro* experiments demonstrated that 7-DHC-derived oxysterols, including 7-KC, caused toxicity to mouse cone, monkey RPE and rat Müller glial cells (Pfeffer et al., 2016). The cause of retinal degeneration by 7-DHC-derived oxysterols in AY9944-treated rats must at least be partially due to oxidative damage, given that co-treatment with

cholesterol and antioxidants is more effective in protecting against retinal degeneration in AY9944-treated rats than treatment of rats with cholesterol alone (Fliesler et al., 2018).

5 OXYSTEROLS AND AGE RELATED MACULAR DEGENERATION

Age-related macular degeneration (AMD) is a chronic disease that affects the macula, the central cone dominated feature of the retina (Mitchell et al., 2018). The disease affects an ageing demographic and inevitably causes a loss of central vision due to growing abnormalities within the macula. AMD is one of the leading causes of visual loss in the developed world, particularly in those of Caucasian ethnicity (Mitchell et al., 2018). This degenerative condition is becoming increasingly prevalent due to an ageing population with predictions suggesting that AMD will double in frequency over the next two decades (Robman et al., 2017). AMD is a complex, multifactorial disease process. Aging is known to be the single most significant risk factor, but both genetic and environmental factors also contribute significantly to AMD etiology. When AMD has progressed to its late stage, it is classified into two major forms: dry (geographic atrophy) and wet (choroidal neovascularisation, CNV). The wet type is the significantly less common (about 10%) of the two, but its progression is fast and severe, accounting for 90% registration of legal blindness caused by AMD. While there is no effective treatment for dry AMD, anti-VEGF therapy is effective in wet AMD, causing regression of CNV and preservation of visual function (Mitchell et al., 2018).

The clinical hallmark of AMD is abnormal accumulation beneath the RPE of extracellular deposits (drusen) and basal linear deposits (BLinD) (Pikuleva and Curcio, 2014). Histopathological analyses has shown that cholesterol is enriched in these deposits, suggesting that defective RCT may play an important role in the pathogenesis of AMD (Curcio et al., 2001, 2005; Wang et al., 2010). RPE cells progressively degenerate in AMD, leading to the death of photoreceptors in the macula. The RPE cells are responsible for cholesterol homeostasis in the retina by supplying cholesterol to photoreceptor cells and by clearing excess cholesterol into the choriocapillaries via the RCT (Biswas et al., 2017; Lyssenko et al., 2018; Storti et al., 2017). Oxysterol-producing genes are abundantly expressed in the RPE (Heo et al., 2011; Liao et al., 2011; Zheng et al., 2012), which is exposed to a

high level of oxidative stress (Datta et al., 2017). Oxidative stress is proposed to induce cholesterol oxidation, so oxysterols may be actively produced in the RPE via enzymatic and free radical reactions. In fact, oxysterol species have been detected and localized to the RPE (Figure 2B) and there may be a functional role for 7-KC, 24S-HC and 27-HC in AMD.

The extremely high levels of 7-KC in the aged RPE and drusen, coupled with an observed increase of 7-KC in the RPE as a result of photodamage, support the notion of a link between 7-KC and AMD (Moreira et al., 2009; Rodriguez and Fliesler, 2009; Rodriguez et al., 2014). Early work showed that 7-KC treatment decreased cell viability, induced cell death and increased ROS production in human RPE cells (Ong et al., 2003; Dugas et al., 2010; Olivier et al., 2016; Rodriguez et al., 2004) and in porcine RPE cells (Joffre et al., 2007). 7-KC is the most abundant oxysterol in oxidized LDL (ox-LDL) (Brown et al., 1996; Oh et al., 2016). Ox-LDL induces inflammation in the RPE and choroidal endothelial cells (Biswas et al., 2017, 2018). 7-KC treatment also induced production of inflammatory cytokines including IL-1 β , IL-6, IL-8, IL-18, TNF- α in RPE cells (Dugas et al., 2010; Huang et al., 2012; Joffre et al., 2007; Larrayoz et al., 2010; Moreira et al., 2009; Shi et al., 2015; Yang et al., 2019), possibly via MAPK/ERK, Akt/PKB, PKC and NF- κ B pathways (Larrayoz et al., 2010; Moreira et al., 2009; Yang et al., 2019). Huang et al (2012) reported that 7-KC also caused endoplasmic reticulum (ER) stress in human RPE cells. Given that the RPE is responsible for the daily renewal of the photoreceptor outer segment via phagocytosis, impaired or decreased phagocytosis is likely to contribute to AMD (Kevany and Palzewski, 2010). Recently, it has been shown that intravitreal injection of 7-KC in rats has resulted in disorganised and decreased apical microvilli of RPE and detachment of apical microvilli from the photoreceptor outer segments, leading to defects in outer segment phagocytosis (Yang et al., 2019).

Retinal microglia are resident macrophages that are distributed in the inner retina of young, healthy individuals. With aging or under stress conditions, these microglia migrate to the outer retina and become activated, demonstrating proinflammatory and proangiogenic properties. Microglial activation is associated with various types of retinal degeneration, including AMD

(Karlstetter et al., 2015). 7-KC has been reported to co-localize with subretinal microglia in aged mice. 7-KC chemoattractively influenced microglial migration to the subretinal space and enhanced microglial activation, indicated by inflammasome activation, increased expression of proinflammatory cytokines (IL-1 β , IL-6, IL-18 and TNF α) and of proangiogenic factors (VEGF, PDGFb and ICAM). Conditional culture media from 7-KC-exposed microglia stimulated endothelial cell migration, while transplantation of 7-KC-exposed microglia in a Matrigel-induced CNV model significantly promoted CNV formation (Indaram et al., 2015).

24S-HC can directly cross the blood-brain barrier and be transported to the liver by circulating lipoproteins, facilitating cholesterol elimination from the brain (Björkhem, 2006). 24S-HC also functions as an activator for nuclear transcription factors, such as LXR α and β (Lehmann et al., 1997), to upregulate expression of cholesterol transport genes including *ABCA1* in both glial cells and neurons (Fukumoto et al., 2002) and *APOE* in astrocytes (Liang et al., 2004), leading to increased cholesterol efflux and mediation of cholesterol homeostasis. Additionally, 24S-HC can suppress Amyloid beta (A β) production by hindering amyloid precursor protein trafficking in Alzheimer's models (Urano et al., 2013). A β has been localized to the drusen of AMD patients and is thought to contribute to AMD pathogenesis (Isas et al., 2010; Luibl et al., 2006). Anti-amyloid therapy suppresses retinal damage and retards vision loss in an AMD mouse model (Ding et al., 2011). Although 24S-HC is reported to be higher in the plasma of AMD patients than that of controls (Lin et al., 2018), the level of 24S-HC in the retina and RPE of AMD patients may be different. So we propose that 24S-HC may play a protective role in AMD by enhancing cholesterol transport and reducing A β formation in neural retina and the RPE.

27-HC is undetectable in neural retina and RPE, possibly due to being immediately oxidized to 27-COOH by CYP27A1 (Mast et al., 2011). Although there is no difference in plasma 27-HC levels of AMD patients and controls (Lin et al., 2018), we believe there to be a connection between 27-HC and AMD based on our recent work (Biswas et al., 2017, 2018). The translocator protein TSPO is a mitochondrial membrane protein that is responsible for delivering cholesterol from the

mitochondrial outer membrane to the inner membrane where it is metabolized to 27-HC by CYP27A1. In turn, 27-HC activates LXR signalling to upregulate the expression of ABCA1, ABCG1/4 and APOE and promotes cholesterol transport in non-steroidogenic cells, including the RPE (Figure 3) (Biswas et al., 2017; Papadopoulos et al., 2015). We found that TSPO is expressed in the human RPE and monkey choroidal endothelial cells. In mouse retina, its expression was extremely high in the RPE and was seen to decline with age. Knockout of TSPO resulted in decreased cholesterol efflux and cholesterol accumulation in the RPE cells. TSPO ligands (FGIN-1-27, [XBD173](#) and [Etifoxine](#)) promoted cholesterol efflux in the RPE and choroidal endothelial cells and upregulated expression of oxysterol metabolism and cholesterol trafficking genes; TSPO ligands also suppressed oxidized LDL-induced oxidative stress and inflammation (Biswas et al., 2017, 2018). The data suggests that TSPO-mediated cholesterol trafficking via 27-HC may play an important role in the pathogenesis of AMD.

7 OXYSTEROLS AND GLAUCOMA

Glaucoma is a leading cause of irreversible blindness; currently affecting nearly 80 million individuals worldwide and predicted to increase to 111.8 million by 2040 (Tham et al., 2014). Glaucoma is characterised by the gradual degeneration of the optic nerve and retinal ganglion cell (RGC) loss (Almasieh et al., 2012; Weinreb et al., 2014). As RGCs are non-dividing cells without the capacity for regeneration, the damage of the optic nerve appears to be irreparable. Loss of RGCs is a complex process, associated with deprivation of neurotrophic factors, failure of axonal transport, mitochondrial dysfunction, neurotoxicity (oxidative stress, inflammation and excitotoxic damage) and defects in RGC-glia interaction (Almasieh et al., 2012). Glaucoma is classified in two main types: primary open-angle glaucoma (POAG) and primary angle-closure glaucoma (PACG). Most cases are POAG and involve a long-term increase in intraocular pressure (IOP). Glaucoma is a complex disease associated with multiple risk factors such as age, genetic background and hyperlipidemia (Wang and Bo, 2019; Weinreb et al., 2014). IOP is considered to be the most common risk factor for glaucoma, and reducing IOP is a well-established clinical approach to slowing glaucomatous degeneration. Although reducing IOP is therapeutically effective in retarding glaucoma progression, there is some

evidence that disease progression is still inevitable (Heijl et al., 2002; Kass et al., 2002), suggesting that neuroprotectants are urgently needed to prevent the degeneration of RGC and optic nerve.

The link between 24S-HC and glaucoma was first proposed when a polymorphism (rs754203) in the intron 2 of *CYP46A1* gene was shown to be associated with increased risk for POAG (Fourgeux et al., 2009). Further work found that elevated IOP in the rat caused a transient increase in expression of CYP46A1 but no change in the retinal 24S-HC between experimental and contralateral eyes (Fourgeux et al., 2012). Inhibition of CYP46A1 in rats by intraperitoneal injection with voriconazole, which has been shown to effectively inhibit CYP46A1 *in vitro* and *in vivo* (Shafaati et al., 2010), led to a 37% decrease in retinal 24S-HC and impaired inner retinal function (ERG test); however, the effect of CYP46A1 inhibition on GCL was not examined (Fourgeux et al., 2014). Ishikawa et al. (2016) found elevated IOP significantly enhanced CYP46A1 expression at mRNA and protein levels in an *ex vivo* rat glaucoma model, while increased CYP46A1 protein was predominantly localized to ganglion cells. Retinal 24S-HC was also increased under conditions of high IOP. The authors also demonstrated that high pressure (75 mmHg) caused damage to the RGCs and that 24S-HC counteracted the damaging effects in a dose-dependent manner. 24S-HC also protected the RGCs from voriconazole-induced damage (Ishikawa et al., 2016). The data suggest that 24S-HC may have a protective role in conditions of high-IOP that are associated with glaucoma.

8 OXYSTEROL-TARGETED THERAPEUTIC STRATEGIES FOR RETINAL DEGENERATION

8.1 Small chemical therapy

The development of oxysterol-targeted therapy will depend on the functional properties of individual oxysterols. For example, when targeting 7-KC, a strategy is needed to inhibit 7-KC-induced toxicity (e.g. oxidative damage and inflammation) and/or to convert 7-KC to less- or non-toxic metabolites; for 24S-HC and 27-HC, ideally there should be more *in situ* biosynthesis. Sterculic acid has been shown to inhibit 7-KC-induced cell death and inflammation in RPE cells and suppressed laser-injury-induced CNV formation in rats (Huang et al., 2012). [Resveratrol](#), a natural polyphenol, has also been shown to reduce 7-KC-induced RPE apoptosis and VEGF secretion (Dugas et al., 2010).

There are other candidates that are worth investigating further. Efavirenz, an anti-HIV drug, has been shown to activate CYP46A1 and improve behaviour in the 5XFAD Alzheimer's disease mouse model (Mast et al., 2017; Petrov et al., 2019). It would be useful to assess the therapeutic potential of Efavirenz via 24S-HC function in both AMD and glaucoma models. TSPO ligands, such as XBD173, promote mitochondrial cholesterol trafficking and increase the metabolism of cholesterol to 27-HC, which upregulates cholesterol transport and metabolism genes via activation of the LXR signal pathway, consequently enhancing intracellular cholesterol removal (Papadopoulos et al., 2015). TSPO ligands have been shown to suppress oxidized LDL-induced oxidative stress and inflammation in the RPE and choroidal endothelial cells (Biswas et al., 2017, 2018). XBD173 can also inhibit microglial activation and attenuate light-induced retinal degeneration (Scholz et al., 2015). Evaluation of the protective effects of XBD173 against 7-KC-induced oxidative damage, inflammation and microglia activation may help to develop new treatments for AMD.

8.2 Gene therapy

Gene therapy is one possible means by which to increase levels of oxysterol-producing enzymes in the retina. Currently, this appears to be a safe and feasible approach for the treatment of inherited retinal degeneration. Adeno-associated virus (AAV) is a commonly used vector for delivery, targeting genes in the retinal cells such as photoreceptors and RPE cells (Lee et al., 2019). Hudry et al (2010) over-expressed CYP46A1 via AAV-mediated gene therapy in the cortex and hippocampus of Alzheimer's disease mice and found that the level of 24S-HC in the brain was significantly increased, while formation of A β deposits and A β -associated pathology were markedly reduced (Hudry et al., 2010). AAV-mediated overexpression of CYP46A1 in the striatum globally regulates cholesterol metabolism and improves behavioural and neuronal functions in Huntington's disease mouse model (Kacher et al., 2019). Similarly, overexpression of CYP46A1 in ganglion cells via gene delivery may help to prevent glaucoma. CYP27A1 is the main enzyme responsible for elimination of 7-KC in the RPE (Heo et al., 2011); it also metabolizes cholesterol to 27-HC, which promotes cholesterol efflux in

the RPE (Biswas et al., 2017). Potentially, then, overexpression of CYP27A1 in the RPE using gene therapy could be a new therapeutic strategy for AMD.

9 CONCLUSIONS

Enriched expression of oxysterol-producing enzymes and the distribution of oxysterols in the retina suggest that oxysterols may have multiple functions in the pathogenesis of degenerative retinal diseases. This review summarizes our current understanding of oxysterol biology in the retina, including biosynthesis, metabolism, function and association with retinal degeneration. Current preclinical and clinical data suggest that 7-KC, 24S-HC and 27-HC are the major oxysterols associated with retinal degeneration. More effort is required to establish a deeper understanding of oxysterol-related pathophysiology in retinal degeneration and to develop new therapeutic interventions to improve visual function in oxysterol-associated retinal disorders.

9.1 Nomenclature of targets and ligands

Key protein targets and ligands in this article are hyperlinked to corresponding entries in <http://www.guidetopharmacology.org>, the common portal for data from the IUPHAR/BPS Guide to PHARMACOLOGY (Harding et al., 2018), and are permanently archived in the Concise Guide to PHARMACOLOGY 2019/20 (Alexander et al., 2019a,b,c).

ACKNOWLEDGEMENT

Research work in Dr Shu's lab was supported by the Rosetrees Trust (M160, M160-F1, M160-F2), National Eye Research Centre (SAC037) and Tenovus Scotland (S20-02). The work was partially supported by the Lotus Scholarship Program of Hunan Province, P. R. China (2019). X.S. is a visiting Professor to Shaoyang University. We are sorry we could not cite other related references.

CONFLICT OF INTEREST

The authors do not have conflicts of interest to declare.

Author Contributions

X.S. conceived this project. X.Z., R.H.A, X.Z., J.R., Z.Z. and X.S. prepared the first draft. X.Z., X.S., and N.S. revised the manuscript.

REFERENCES

- Adams, C.M., Reitz J, DeBrabander JK, Feramisco, J.D., Brown, M.S., & Goldstein, J.L. (2004). Cholesterol and 25-Hydroxycholesterol Inhibit Activation of SREBPs by Different Mechanisms, Both Involving SCAP and Insigs. *Journal of Biological Chemistry*, 279, 52772–52780.
- Alexander, S. P., Cidlowski, J. A., Kelly, E., Mathie, A., Peters, J. A., Veale, E. L., ... Sharman, J. L. (2019a). The Concise Guide to PHARMACOLOGY 2019/20: Nuclear hormone receptors. *British Journal of Pharmacology*, 176, S229–S246. <https://doi.org/10.1111/bph.14750>
- Alexander, S.P., Fabbro, D., Kelly, E., Mathie, A., Peters, J.A., Veale, E.L., Armstrong, J.F., Faccenda, E., Harding, S.D., Pawson, A.J. and Sharman, J.L. (2019b). The concise guide to pharmacology 2019/20: Enzymes. *British Journal of Pharmacology*, 176, S297-S396.
- Alexander, S. P., Kelly, E., Mathie, A., Peters, J. A., Veale, E. L., Armstrong, J. F., ... Southan, C. (2019c). The Concise Guide to PHARMACOLOGY 2019/20: Transporters. *British Journal of Pharmacology*, 176, S397–S493. <https://doi.org/10.1111/bph.14753>
- Almasieh, M., Wilson, A.M., Morquette, B., Vargas, J.L.C., & Di Polo, A. (2012). The molecular basis of retinal ganglion cell death in glaucoma. *Progress in retinal and eye research*, 31(2), 152-181.
- Anderson, A., Campo, A., Fulton, E., Corwin, A., Jerome III, W.G., & O'Connor, M.S. (2020). 7-Ketocholesterol in disease and aging. *Redox Biology*, 29, 101380.
- Atchaneeyasakul, L.O., Linck, L.M., Connor, W.E., Weleber, R.G., & Steiner, R.D. (1998). Eye findings in 8 children and a spontaneously aborted fetus with RSH/Smith-Lemli-Opitz syndrome. *American journal of medical genetics*, 80(5), 501-505.
- Biswas, L., Farhan, F., Reilly, J., Bartholomew, C., & Shu, X. (2018). TSPO ligands promote cholesterol efflux and suppress oxidative stress and inflammation in choroidal endothelial cells. *International journal of molecular sciences*, 19(12), 3740.
- Biswas, L., Zhou, X., Dhillon, B., Graham, A., & Shu, X. (2017) Retinal pigment epithelium cholesterol efflux mediated by the 18kDa translocator protein, TSPO, a potential target for treating age-related macular degeneration. *Human Molecular Genetics*, 26, 4327–4339.

571 Björkhem, I. (2006). Crossing the barrier: oxysterols as cholesterol transporters and metabolic
572 modulators in the brain. *Journal of internal medicine*, 260(6), 493-508.

573 Bodin, K., Andersson, U., Rystedt, E., Ellis, E., Norlin, M., Pikuleva, I., Eggertsen, G., Björkhem, I. and
574 Diczfalusy, U., 2002. Metabolism of 4 β -hydroxycholesterol in humans. *Journal of Biological*
575 *Chemistry*, 277(35), 31534-31540.

576 Bretilon, L., Thuret, G., Grégoire, S., Acar, N., Joffre, C., Bron, A.M., ... Creuzot-Garcher, C.P. (2008).
577 Lipid and fatty acid profile of the retina, retinal pigment epithelium/choroid, and the lacrimal gland,
578 and associations with adipose tissue fatty acids in human subjects. *Experimental eye research*, 87(6),
579 521-528.

580 Bretilon, L., Diczfalusy, U., Björkhem, I., Maire, M.A., Martine, L., Joffre, C., ... Creuzot-Garcher, C.
581 (2007). Cholesterol-24S-hydroxylase (CYP46A1) is specifically expressed in neurons of the neural
582 retina. *Current eye research*, 32(4), 361-366.

583 Brown, A.J., Dean, R.T. and Jessup, W., 1996. Free and esterified oxysterol: formation during copper-
584 oxidation of low density lipoprotein and uptake by macrophages. *Journal of lipid research*, 37(2),
585 320-335.

586 Cruysberg, J.R.M., Wevers, R.A., Van Engelen, B.G.M., Pinckers, A.J.L.G., van Spreeken, A.C.G.A., &
587 Tolboom, J.J.M. (1995). Ocular and systemic manifestations of cerebrotendinous
588 xanthomatosis. *American journal of ophthalmology*, 120(5), 597-604.

589 Curcio, C.A., Millican, C.L., Bailey, T., & Kruth, H.S. (2001). Accumulation of cholesterol with age in
590 human Bruch's membrane. *Investigative ophthalmology & visual science*, 42(1), 265-274.

591 Curcio, C.A., Presley, J.B., Malek, G., Medeiros, N.E., Avery, D.V., & Kruth, H.S. (2005). Esterified and
592 unesterified cholesterol in drusen and basal deposits of eyes with age-related
593 maculopathy. *Experimental eye research*, 81(6), 731-741.

594 Datta, S., Cano, M., Ebrahimi, K., Wang, L., & Handa, J.T. (2017). The impact of oxidative stress and
595 inflammation on RPE degeneration in non-neovascular AMD. *Progress in retinal and eye research*, 60,
596 201-218.

597 DeBose-Boyd RA, Brown MS, Li W-P, Nohturfft A, Goldstein JL, Espenshade PJ. 1999. Transport
 598 dependent proteolysis of SREBP: Relocation of site-1 protease from Golgi to ER obviates the need for
 599 SREBP transport to Golgi. *Cell* 99:703–12
 600 Dotti, M.T., Rufa, A., & Federico, A. (2001). Cerebrotendinous xanthomatosis: heterogeneity of
 601 clinical phenotype with evidence of previously undescribed ophthalmological findings. *Journal of*
 602 *inherited metabolic disease*, 24(7), 696-706.
 603 Dubrac, S., Lear, S.R., Ananthanarayanan, M., Balasubramaniyan, N., Bollineni, J., Shefer, S., ... Batta,
 604 A.K. (2005). Role of CYP27A in cholesterol and bile acid metabolism. *Journal of lipid research*, 46(1),
 605 76-85.
 606 Dugas, B., Charbonnier, S., Baarine, M., Ragot, K., Delmas, D., Ménétrier, F., ... Creuzot-Garcher, C.
 607 (2010). Effects of oxysterols on cell viability, inflammatory cytokines, VEGF, and reactive oxygen
 608 species production on human retinal cells: cytoprotective effects and prevention of VEGF secretion
 609 by resveratrol. *European journal of nutrition*, 49(7), 435-446.
 610 El-Darzi, N., Astafev, A., Mast, N., Saadane, A., Lam, M., & Pikuleva, I.A. (2018). N, N-Dimethyl-3β-
 611 hydroxycholeamide reduces retinal cholesterol via partial inhibition of retinal cholesterol
 612 biosynthesis rather than its liver X receptor transcriptional activity. *Frontiers in pharmacology*, 9, 827.
 613 Elias, E.R., Hansen, R.M., Irons, M., Quinn, N.B., & Fulton, A.B. (2003). Rod photoreceptor responses
 614 in children with Smith-Lemli-Opitz syndrome. *Archives of Ophthalmology*, 121(12), 1738-1743.
 615 Escher, G., Krozowski, Z., Croft, K.D. & Sviridov, D. (2003). Expression of Sterol 27-Hydroxylase
 616 (CYP27A1) Enhances Cholesterol Efflux. *Journal of Biological Chemistry*, 278, 11015-11019.
 617 Fitzky, B.U., Moebius, F.F., Asaoka, H., Waage-Baudet, H., Xu, L., Xu, G., ... Batta, A.K. (2001). 7-
 618 Dehydrocholesterol-dependent proteolysis of HMG-CoA reductase suppresses sterol biosynthesis in
 619 a mouse model of Smith-Lemli-Opitz/RSH syndrome. *The Journal of clinical investigation*, 108(6),
 620 905-915.
 621 Fliesler, S. J., & Bretillon, L. (2010). The ins and outs of cholesterol in the vertebrate retina. *Journal of*
 622 *lipid research*, 51(12), 3399-3413.

623 Fliesler, S.J., Peachey, N.S., Richards, M.J., Nagel, B.A., & Vaughan, D.K. (2004). Retinal degeneration
624 in a rodent model of Smith-Lemli-Opitz syndrome: electrophysiologic, biochemical, and morphologic
625 features. *Archives of Ophthalmology*, 122(8), 1190-1200.

626 Fliesler, S.J., Peachey, N.S., Herron, J., Hines, K.M., Weinstock, N.I., Rao, S.R., & Xu, L. (2018).
627 Prevention of retinal degeneration in a rat model of Smith-Lemli-Opitz syndrome. *Scientific*
628 *reports*, 8(1), 1-13.

629 Fliesler, S.J., Richards, M.J., Miller, C.Y., Peachey, N.S. (1999). Marked alteration of sterol metabolism
630 and composition without compromising retinal development or function. *Investigative*
631 *ophthalmology & visual science*, 40(8), 1792-1801.

632 Fliesler, S.J., & Xu, L. (2018). Oxysterols and Retinal Degeneration in a Rat Model of Smith-Lemli-
633 Opitz Syndrome: Implications for an Improved Therapeutic Intervention. *Molecules*, 23(10), 2720.

634 Fourgeux, C., Martine, L., Acar, N., Bron, A.M., Creuzot-Garcher, C.P., & Bretillon, L. (2014). In vivo
635 consequences of cholesterol-24S-hydroxylase (CYP46A1) inhibition by voriconazole on cholesterol
636 homeostasis and function in the rat retina. *Biochemical and biophysical research*
637 *communications*, 446(3), 775-781.

638 Fourgeux, C., Martine, L., Björkhem, I., Diczfalusy, U., Joffre, C., Acar, N., ... Bretillon, L. (2009).
639 Primary open-angle glaucoma: association with cholesterol 24S-hydroxylase (CYP46A1) gene
640 polymorphism and plasma 24-hydroxycholesterol levels. *Investigative ophthalmology & visual*
641 *science*, 50(12), 5712-5717.

642 Fourgeux, C., Martine, L., Pasquis, B., Maire, M.A., Acar, N., Creuzot-Garcher, C., ... Bretillon, L.
643 (2012). Steady-state levels of retinal 24S-hydroxycholesterol are maintained by glial cells
644 intervention after elevation of intraocular pressure in the rat. *Acta ophthalmologica*, 90(7), e560-
645 e567.

646 Fu, X., Menke, J.G., Chen, Y., Zhou, G., MacNaul, K.L., Wright, S.D., Sparrow, C.P. and Lund, E.G., 2001.
647 27-hydroxycholesterol is an endogenous ligand for liver X receptor in cholesterol-loaded
648 cells. *Journal of Biological Chemistry*, 276(42), 38378-38387.

649 Fukumoto, H., Deng, A., Irizarry, M.C., Fitzgerald, M.L., & Rebeck, G.W. (2002). Induction of the
650 cholesterol transporter ABCA1 in central nervous system cells by liver x receptor agonists increases
651 secreted abeta levels. *J Biol Chem* 277:48508–48513

652 Garry, D., Hansen, R.M., Moskowitz, A., Elias, E.R., Irons, M., Fulton, A.B. (2010). Cone ERG responses
653 in patients with Smith-Lemli-Opitz syndrome (SLOS). *Documenta ophthalmologica*, 121(2), 85-91.

654 Griffiths, W.J., & Wang, Y. (2019). Oxysterol research: a brief review. *Biochemical Society*
655 *Transactions*, 47(2), 517-526.

656 Guarneri, P., Guarneri, R., Cascio, C., Pavasant, P., Piccoli, F., & Papadopoulos, V. (1994)
657 Neurosteroidogenesis in rat retinas. *Journal of Neurochemistry*, 63, 86–96.

658 Guillemot-Legris, O., Mutemberezi, V., & Muccioli, G.G. (2016). Oxysterols in metabolic syndrome:
659 from bystander molecules to bioactive lipids. *Trends in molecular medicine*, 22(7), 594-614.

660 Harding, S.D., Sharman, J.L., Faccenda, E., Southan, C., Pawson AJ, Ireland S, Gray AJG, Bruce L,
661 Alexander SPH, Anderton S, Bryant C, Davenport AP, Doerig C, Fabbro D, Levi-Schaffer F, Spedding M,
662 Davies JA, NC-IUPHAR (2018). The IUPHAR/BPS Guide to PHARMACOLOGY in 2018: updates and
663 expansion to encompass the new guide to IMMUNOPHARMACOLOGY. *Nucleic Acids Res.* 46, D1091-
664 1106. doi:10.1093/nar/gkx1121.

665 Heijl, A., Leske, M.C., Bengtsson, B., Hyman, L., Bengtsson, B., & Hussein, M. (2002). Reduction of
666 intraocular pressure and glaucoma progression: results from the Early Manifest Glaucoma
667 Trial. *Archives of ophthalmology*, 120(10), 1268-1279.

668 Heo, G.Y., Bederman, I., Mast, N., Liao, W.L., Turko, I.V., & Pikuleva, I.A. (2011). Conversion of 7-
669 ketocholesterol to oxysterol metabolites by recombinant CYP27A1 and retinal pigment epithelial
670 cells. *Journal of lipid research*, 52(6), 1117-1127.

671 Honda, A., Salen, G., Matsuzaki, Y., Batta, A.K., Xu, G., Leitersdorf, E., ... Shefer, S. (2001). Differences
672 in hepatic levels of intermediates in bile acid biosynthesis between Cyp27^{-/-} mice and CTX. *Journal*
673 *of Lipid Research*, 42(2), 291-300.

674 Hu, D.N., Simon, J.D., & Sarna, T. (2008). Role of ocular melanin in ophthalmic physiology and
675 pathology. *Photochemistry and photobiology*, 84(3), 639-644.

676 Huang, J.D., Amaral, J., Lee, J.W., Larrayoz, I.M., & Rodriguez, I.R. (2012). Sterculic acid antagonizes
677 7-ketocholesterol-mediated inflammation and inhibits choroidal neovascularization. *Biochimica et*
678 *Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids*, 1821(4), 637-646.

679 Hua, X., Yokoyama, C., Wu, J., Briggs, M.R., Brown, M.S, Goldstein, J.L., ... Wang, X. (1993). SREBP-2, a
680 second basic-helix-loop-helix-leucine zipper protein that stimulates transcription by binding to a
681 sterol regulatory element. *Proceedings of the National Academy of Sciences of the United States of*
682 *America*, 90, 11603-7

683 Hudry, E., Van Dam, D., Kulik, W., De Deyn, P.P., Stet, F.S., Ahouansou, O., ... Cartier, N. (2010).
684 Adeno-associated virus gene therapy with cholesterol 24-hydroxylase reduces the amyloid pathology
685 before or after the onset of amyloid plaques in mouse models of Alzheimer's disease. *Molecular*
686 *Therapy*, 18(1), 44-53.

687 Indaram, M., Ma, W., Zhao, L., Fariss, R.N., Rodriguez, I.R., & Wong, W.T. (2015). 7-Ketocholesterol
688 increases retinal microglial migration, activation, and angiogenicity: a potential pathogenic
689 mechanism underlying age-related macular degeneration. *Scientific reports*, 5, 9144.

690 Ishikawa, M., Yoshitomi, T., Covey, D.F., Zorumski, C.F., & Izumi, Y. (2018). Neurosteroids and
691 oxysterols as potential therapeutic agents for glaucoma and Alzheimer's
692 disease. *Neuropsychiatry*, 8(1), 344.

693 Isas, J.M., Luibl, V., Johnson, L.V., Kaye, R., Wetzel, R., Glabe, C.G., ..., Jeannie Chen, J.
694 (2010). Soluble and mature amyloid fibrils in drusen deposits. *Investigative ophthalmology & visual*
695 *science*, 51:1304-1310.

696 Ishikawa, M., Yoshitomi, T., Zorumski, C.F., & Izumi, Y. (2016). 24 (S)-Hydroxycholesterol protects the
697 ex vivo rat retina from injury by elevated hydrostatic pressure. *Scientific reports*, 6, 33886.

698 Jaliffa, C.O., Howard, S., Hoijman, E., Salido, E., Sarmiento, M.I.K., Arias, P., & Rosenstein, R.E. (2005).
 699 Effect of neurosteroids on the retinal gabaergic system and electroretinographic activity in the
 700 golden hamster. *Journal of neurochemistry*, 94(6), 1666-1675.

701 Janowski, B.A., Grogan, M.J., Jones, S.A., Wisely, G.B., Kliewer, S.A., Corey, E.J. and Mangelsdorf, D.J.,
 702 1999. Structural requirements of ligands for the oxysterol liver X receptors LXR α and
 703 LXR β . *Proceedings of the National Academy of Sciences*, 96(1), 266-271.

704 Joffre, C., Leclère, L., Buteau, B., Martine, L., Cabaret, S., Malvitte, L., ... Bretillon, L. (2007).
 705 Oxysterols induced inflammation and oxidation in primary porcine retinal pigment epithelial
 706 cells. *Current eye research*, 32(3), 271-280.

707 Karlstetter, M., Scholz, R., Rutar, M., Wong, W.T., Provis, J.M., & Langmann, T. (2015). Retinal
 708 microglia: just bystander or target for therapy?. *Progress in retinal and eye research*, 45, 30-57.

709 Kass, M.A., Heuer, D.K., Higginbotham, E.J., Johnson, C.A., Keltner, J.L., Miller, J.P., ... Gordon, M.O.,
 710 2002. The Ocular Hypertension Treatment Study: a randomized trial determines that topical ocular
 711 hypotensive medication delays or prevents the onset of primary open-angle glaucoma. *Archives of*
 712 *ophthalmology*, 120(6), 701-713.

713 Katz, M.L., & Gao, C.L. (1995). Vitamin A incorporation into lipofuscin-like inclusions in the retinal
 714 pigment epithelium. *Mechanisms of ageing and development*, 84(1), 29-38.

715 Kevany, B.M., & Palczewski, K. (2010). Phagocytosis of retinal rod and cone
 716 photoreceptors. *Physiology*, 25(1), 8-15.

717 Kolf-Clauw, M., Chevy, F., Wolf, C., Siliart, B., Citadelle, D., & Roux, C. (1996). Inhibition of 7-
 718 dehydrocholesterol reductase by the teratogen AY9944: A rat model for Smith-Lemli-Opitz
 719 syndrome. *Teratology*, 54(3), 115-125.

720 Korade, Z., Xu, L., Shelton, R., & Porter, N.A. (2010). Biological activities of 7-dehydrocholesterol-
 721 derived oxysterols: implications for Smith-Lemli-Opitz syndrome. *Journal of lipid research*, 51(11),
 722 3259-3269.

723 Kretzer, F.L., Hittner, H.M., & Mehta, R.S. (1981). Ocular manifestations of the Smith-Lemli-Opitz
724 syndrome. *Archives of Ophthalmology*, 99(11), 2000-2006.

725 Lamb, T.D., Collin, S.P., & Pugh Jr, E.N. (2007). Evolution of the vertebrate eye: opsins,
726 photoreceptors, retina and eye cup. *Nature Reviews Neuroscience*, 8(12), 960-976

727 Larrayoz, I.M., Huang, J.D., Lee, J.W., Pascual, I., & Rodríguez, I.R. (2010). 7-Ketocholesterol-induced
728 inflammation: involvement of multiple kinase signaling pathways via NFκB but independently of
729 reactive oxygen species formation. *Investigative ophthalmology & visual science*, 51(10), 4942-4955.

730 Liang Y, Lin S, Beyer TP, Zhang Y, Wu X, Bales KR, DeMattos RB, May PC, Li SD, Jiang XC Eacho PI, Cao
731 G, Paul SM (2004) A liver X receptor and retinoid X receptor heterodimer mediates apolipoprotein E
732 expression secretion and cholesterol homeostasis in astrocyte. *J NeuroChem* 88:623–634

733 Lee, J.W., Fuda, H., Javitt, N.B., Strott, C.A., & Rodriguez, I.R. (2006). Expression and localization of
734 sterol 27-hydroxylase (CYP27A1) in monkey retina. *Experimental eye research*, 83(2), 465-469.

735 Lee, J.H., Wang, J.H., Chen, J., Li, F., Edwards, T.L., Hewitt, A.W., & Liu, G.S. (2019). Gene therapy for
736 visual loss: Opportunities and concerns. *Progress in retinal and eye research*, 68, 31-53.

737 Lehmann JM, Kliewer SA, Moore LB, Smith-Oliver TA, Oliver BB, Su JL, Sundseth SS, Winegar
738 DA, Blanchard DE, Spencer TA, Willson TM. (1997). Activation of the nuclear receptor LXR by
739 oxysterols defines a new hormone response pathway. *The Journal of Biological Chemistry*, 272(6),
740 3137–3140.

741 Liao, W.L., Heo, G.Y., Dodder, N.G., Reem, R.E., Mast, N., Huang, S., ... Pikuleva, I.A. (2011).
742 Quantification of Cholesterol-Metabolizing P450s CYP27A1 and CYP46A1 in Neural Tissues Reveals a
743 Lack of Enzyme– Product Correlations in Human Retina but Not Human Brain. *Journal of proteome*
744 *research*, 10(1), 241-248.

745 Lin, J.B., Sene, A., Santeford, A., Fujiwara, H., Sidhu, R., Ligon, M.M., ... Apte, R.S. (2018). Oxysterol
746 signatures distinguish age-related macular degeneration from physiologic aging. *EBioMedicine*, 32, 9-
747 20.

748 Lin, J.B., Mast, N., Bederman, I.R., Li, Y., Brunengraber, H., Björkhem, I., & Pikuleva, I.A. (2016).
 749 Cholesterol in mouse retina originates primarily from in situ de novo biosynthesis. *Journal of lipid*
 750 *research*, 57(2), 258-264
 751 Luibl, V., Isas, J.M., Kaye, R., Glabe, C.G., Langen, R., Chen, J. (2006) Drusen deposits associated with
 752 aging and age-related macular degeneration contain nonfibrillar amyloid oligomers. *Journal of*
 753 *Clinical Investigation*, 116, 378–385.
 754 Lund, E.G., Xie, C., Kotti, T., Turley, S.D., Dietschy, J.M., & Russell, D.W. (2003). Knockout of the
 755 cholesterol 24-hydroxylase gene in mice reveals a brain-specific mechanism of cholesterol
 756 turnover. *Journal of Biological Chemistry*, 278(25), 22980-22988.
 757 Lyssenko, N.N., Haider, N., Picataggi, A., Cipollari, E., Jiao, W., Phillips, M.C., ... Chavali, V.R.M. (2018).
 758 Directional ABCA1-mediated cholesterol efflux and apoB-lipoprotein secretion in the retinal pigment
 759 epithelium. *Journal of lipid research*, 59(10), 1927-1939.
 760 Mast, N., Reem, R., Bederman, I., Huang, S., DiPatre, P.L., Björkhem, I., & Pikuleva, I.A. (2011).
 761 Cholestenic acid is an important elimination product of cholesterol in the retina: comparison of
 762 retinal cholesterol metabolism with that in the brain. *Investigative ophthalmology & visual*
 763 *science*, 52(1), 594-603.
 764 Mast, N., Saadane, A., Valencia-Olvera, A., Constans, J., Maxfield, E., Arakawa, H., ... Pikuleva, I.A.
 765 (2017). Cholesterol-metabolizing enzyme cytochrome P450 46A1 as a pharmacologic target for
 766 Alzheimer's disease. *Neuropharmacology*, 123, 465-476.
 767 Mazzone, T., Reardon, C. (1994) Expression of heterologous human apolipoprotein E by J774
 768 macrophages enhances cholesterol efflux to HDL3. *Journal of Lipid Research*, 35, 1345–1353.
 769 Mitchell, P., Liew, G., Gopinath, B., & Wong, T.Y. (2018). Age-related macular degeneration. *The*
 770 *Lancet*, 392(10153), 1147-1159.
 771 Moreira, E.F., Larrayoz, I.M., Lee, J.W., & Rodríguez, I.R. (2009). 7-Ketocholesterol is present in lipid
 772 deposits in the primate retina: potential implication in the induction of VEGF and CNV
 773 formation. *Investigative ophthalmology & visual science*, 50(2), 523-532.

774 Morgan, S.J., McKenna, P., & Bosanquet, R.C. (1989). Case of cerebrotendinous xanthomatosis. I:
 775 Unusual ophthalmic features. *British journal of ophthalmology*, 73(12), 1011-1014.

776 Murphy, R.C. and Johnson, K.M. (2008) Cholesterol, reactive oxygen species, and the formation of
 777 biologically active mediators. *J. Biol. Chem.* **283**, 15521–15525

778 Mustafi, D., Engel, A.H., & Palczewski, K. (2009). Structure of cone photoreceptors. *Progress in*
 779 *retinal and eye research*, 28(4), 289-302.

780 Nowaczyk, M.J., & Irons, M.B. (2012), November. Smith–Lemli–Opitz syndrome: phenotype, natural
 781 history, and epidemiology. In *American Journal of Medical Genetics Part C: Seminars in Medical*
 782 *Genetics* (Vol. 160, No. 4, pp. 250-262). Hoboken: Wiley Subscription Services, Inc., A Wiley Company.

783 Oh, M.J., Zhang, C., LeMaster, E., Adamos, C., Berdyshev, E., Bogachkov, Y., Kohler, E.E., Baruah, J.,
 784 Fang, Y., Schraufnagel, D.E. and Wary, K.K., 2016. Oxidized LDL signals through Rho-GTPase to induce
 785 endothelial cell stiffening and promote capillary formation. *Journal of lipid research*, 57(5), 791-808.

786 Olivier, E., Dutot, M., Regazzetti, A., Leguillier, T., Dargere, D., Auzeil, N., ... Rat, P., 2016. P2X7-
 787 pannexin-1 and wmyloid β -induceu oxysterol input in human retinal cell: Role in age-related macular
 788 degeneration?. *Biochimie*, 127, 70-78.

789 Omarova, S., Charvet, C.D., Reem, R.E., Mast, N., Zheng, W., Huang, S., ... Pikuleva, I.A. (2012).
 790 Abnormal vascularization in mouse retina with dysregulated retinal cholesterol homeostasis. *The*
 791 *Journal of clinical investigation*, 122(8), 3012-3023.

792 Ong, J.M., Aoki, A.M., Seigel, G.M., Sacerio, I., Castellon, R., Nesburn, A.B., & Kenney, M.C. (2003).
 793 Oxysterol-induced toxicity in R28 and ARPE-19 cells. *Neurochemical research*, 28(6), 883-891.

794 Papadopoulos, V., Aghazadeh, Y., Fan, J., Campioli, E., Zirkin, B., & Midzak, A. (2015). Translocator
 795 protein-mediated pharmacology of cholesterol transport and steroidogenesis. *Molecular and cellular*
 796 *endocrinology*, 408, 90-98.

797 Petrov, A.M., Astafev, A.A., Mast, N., Saadane, A., El-Darzi, N., & Pikuleva, I.A. (2019). The Interplay
 798 between Retinal Pathways of Cholesterol Output and Its Effects on Mouse
 799 Retina. *Biomolecules*, 9(12), 867.

800 Petrov, A.M., Lam, M., Mast, N., Moon, J., Li, Y., Maxfield, E., & Pikuleva, I.A. (2019). CYP46A1
 801 activation by efavirenz leads to behavioral improvement without significant changes in amyloid
 802 plaque load in the brain of 5XFAD mice. *Neurotherapeutics*, 16(3), 710-724.
 803 Pfeiffer, B.A., Xu, L., Porter, N.A., Rao, S.R., & Fliesler, S.J. (2016). Differential cytotoxic effects of 7-
 804 dehydrocholesterol-derived oxysterols on cultured retina-derived cells: Dependence on sterol
 805 structure, cell type, and density. *Experimental eye research*, 145, 297-316.
 806 Pikuleva, I.A., & Curcio, C.A. (2014). Cholesterol in the retina: the best is yet to come. *Progress in*
 807 *retinal and eye research*, 41, 64-89.
 808 Purves, D., Augustine, G.J., Fitzpatrick, D., Katz, L.C., LaMantia, A.S., McNamara, J.O., & Williams, S.M.
 809 (2001). The retina, Neuroscience. 2nd edition, Sinauer Associates, Inc.
 810 Radhakrishnan A, Yukio Ikeda Y, Kwon HJ, Brown MS, Goldstein JL. Sterol-regulated transport of
 811 SREBPs from endoplasmic reticulum to Golgi: oxysterols block transport by binding to Insig. The
 812 National Academy of Sciences of the USA, 104 (16): 6511-6518
 813 Rodriguez, I.R., Alam, S., & Lee, J.W. (2004). Cytotoxicity of oxidized low-density lipoprotein in
 814 cultured RPE cells is dependent on the formation of 7-ketocholesterol. *Investigative ophthalmology*
 815 *& visual science*, 45(8), 830-2837.
 816 Ramirez, D.M., Andersson, S., & Russell, D.W. (2008). Neuronal expression and subcellular
 817 localization of cholesterol 24-hydroxylase in the mouse brain. *Journal of Comparative*
 818 *Neurology*, 507(5), 1676-1693.
 819 Remington, L.A., Chapter 4 - Retina, Editor(s): Lee Ann Remington, Clinical Anatomy and Physiology
 820 of the Visual System (Third Edition), ButterworthHeinemann, 2012, Pages 61-92,
 821 Repa, J.J., Lund, E.G., Horton, J.D., Leitersdorf, E., Russell, D.W., Dietschy, J.M., & Turley, S.D. (2000).
 822 Disruption of the sterol 27-hydroxylase gene in mice results in hepatomegaly and
 823 hypertriglyceridemia reversal by Cholic acid feeding. *Journal of Biological Chemistry*, 275(50), 39685-
 824 39692.

825 Robman, L., Guymer, R., Woods, R., Ward, S., Wolfe, R., Phung, J., ... Lockery, J. (2017). Age-related
826 macular degeneration in a randomized controlled trial of low-dose aspirin: Rationale and study
827 design of the ASPREE-AMD study. *Contemporary clinical trials communications*, 6, 105-114.

828 Rodriguez, I.R., Alam, S., & Lee, J.W. (2004). Cytotoxicity of oxidized low-density lipoprotein in
829 cultured RPE cells is dependent on the formation of 7-ketocholesterol. *Investigative ophthalmology
830 & visual science*, 45(8), 2830-2837.

831 Rodriguez, I.R., & Fliesler, S.J. (2009). Photodamage generates 7-keto-and 7-hydroxycholesterol in
832 the rat retina via a free radical-mediated mechanism. *Photochemistry and photobiology*, 85(5), 1116-
833 1125.

834 Rodriguez, I.R., Clark, M.E., Lee, J.W., & Curcio, C.A. (2014). 7-ketocholesterol accumulates in ocular
835 tissues as a consequence of aging and is present in high levels in drusen. *Experimental eye
836 research*, 128, 151-155.

837 Rosen, H., Reshef, A., Maeda, N., Lippoldt, A., Shpizen, S., Triger, L., ... Leitersdorf, E. (1998).
838 Markedly reduced bile acid synthesis but maintained levels of cholesterol and vitamin D metabolites
839 in mice with disrupted sterol 27-hydroxylase gene. *Journal of Biological Chemistry*, 273(24), 14805-
840 14812.

841 Saadane, A., Mast, N., Charvet, C.D., Omarova, S., Zheng, W., Huang, S.S., ... Pikuleva, I.A. (2014).
842 Retinal and nonocular abnormalities in Cyp27a1^{-/-} Cyp46a1^{-/-} mice with dysfunctional metabolism
843 of cholesterol. *The American journal of pathology*, 184(9), 2403-2419.

844 Saadane, A., Mast, N., Dao, T., Ahmad, B., & Pikuleva, I.A. (2016). Retinal hypercholesterolemia
845 triggers cholesterol accumulation and esterification in photoreceptor cells. *Journal of Biological
846 Chemistry*, 291(39), 20427-20439.

847 Saadane, A., Mast, N., Trichonas, G., Chakraborty, D., Hammer, S., Busik, J.V., ... Pikuleva, I.A. (2019).
848 Retinal Vascular Abnormalities and Microglia Activation in Mice with Deficiency in Cytochrome P450
849 46A1-Mediated Cholesterol Removal. *The American journal of pathology*, 189(2), 405-425.

850 Shafaati, M., Mast, N., Beck, O., Nayef, R., Heo, G.Y., Björkhem-Bergman, L., ... Pikuleva, I.A. (2010).
851 The antifungal drug voriconazole is an efficient inhibitor of brain cholesterol 24S-hydroxylase in vitro
852 and in vivo. *Journal of lipid research*, 51(2), 318-323.

853 Scholz, R., Caramoy, A., Bhuckory, M.B., Rashid, K., Chen, M., Xu, H., ... Langmann, T. (2015).
854 Targeting translocator protein (18 kDa)(TSPO) dampens pro-inflammatory microglia reactivity in the
855 retina and protects from degeneration. *Journal of neuroinflammation*, 12(1), 201.

856 Shi, G., Chen, S., Wandu, W.S., Ogbeifun, O., Nugent, L.F., Maminishkis, A., ... Gery, I. (2015).
857 Inflammasomes induced by 7-Ketocholesterol And other stimuli in RPE and in bone marrow-derived
858 cells differ markedly in their production of IL-1 β and IL-18. *Investigative ophthalmology & visual*
859 *science*, 56(3), 1658-1664.

860 Solomon, S.G., & Lennie, P. (2007). The machinery of colour vision. *Nature Reviews Neuroscience*,
861 8(4), 276-286.

862 Steinberg, R.H. (1985). Interactions between the retinal pigment epithelium and the neural retina.
863 *Documenta Ophthalmologica*, 60(4), 327-346.

864 Storti, F., Raphael, G., Griesser, V., Klee, K., Drawnel, F., Willburger, C., ... Grimm, C. (2017).
865 Regulated efflux of photoreceptor outer segment-derived cholesterol by human RPE
866 cells. *Experimental eye research*, 165, 65-77.

867 Strauss, O. (2005). The retinal pigment epithelium in visual function. *Physiological reviews*, 85(3),
868 845-881.

869 Tham, Y.C., Li, X., Wong, T.Y., Quigley, H.A., Aung, T., & Cheng, C.Y. (2014). Global prevalence of
870 glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-
871 analysis. *Ophthalmology*, 121(11), 2081-2090.

872 Tserentsoodol, N., Sztejn, J., Campos, M., Gordiyenko, N.V., Fariss, R.N., Lee, J.W., Fliesler,
873 S.J., Rodriguez, I.R. (2006). Uptake of cholesterol by the retina occurs primarily via a low density
874 lipoprotein receptor-mediated process. *Molecular Vision*, 12, 1306e 1318.

875 Urano, Y., Ochiai, S., & Noguchi, N. (2013). Suppression of amyloid- β production by 24S-
876 hydroxycholesterol via inhibition of intracellular amyloid precursor protein trafficking. *The FASEB*
877 *Journal*, 27(10), 4305-4315.

878 Wang, S., & Bao, X. (2019). Hyperlipidemia, blood lipid level, and the risk of glaucoma: a meta-
879 analysis. *Investigative ophthalmology & visual science*, 60(4), 1028-1043.

880 Wang, L., Clark, M.E., Crossman, D.K., Kojima, K., Messinger, J.D., Mobley, J.A., & Curcio, C.A. (2010).
881 Abundant lipid and protein components of drusen. *PloS one*, 5(4).

882 Wassif, C.A., Zhu, P., Kratz, L., Krakowiak, P.A., Battaile, K.P., Weight, F.F., ... Stewart, R.R. (2001).
883 Biochemical, phenotypic and neurophysiological characterization of a genetic mouse model of
884 RSH/Smith–Lemli–Opitz syndrome. *Human Molecular Genetics*, 10(6), 555-564.

885 Weinreb, R.N., Aung, T., & Medeiros, F.A. (2014). The pathophysiology and treatment of glaucoma: a
886 review. *Jama*, 311(18), 1901-1911.

887 Wright, A.F., Chakarova, C.F., El-Aziz, M.M.A., & Bhattacharya, S.S. (2010). Photoreceptor
888 degeneration: genetic and mechanistic dissection of a complex trait. *Nature Reviews Genetics*, 11(4),
889 273-284.

890 Xu, L., Davis, T.A., & Porter, N.A. (2009). Rate constants for peroxidation of polyunsaturated fatty
891 acids and sterols in solution and in liposomes. *Journal of the American Chemical Society*, 131(36),
892 13037-13044.

893 Xu, L., Korade, Z. and Porter, N.A., 2010. Oxysterols from free radical chain oxidation of 7-
894 dehydrocholesterol: product and mechanistic studies. *Journal of the American Chemical*
895 *Society*, 132(7), 2222-2232.

896 Xu, L., Liu, W., Sheflin, L.G., Fliesler, S.J., & Porter, N.A. (2011). Novel oxysterols observed in tissues
897 and fluids of AY9944-treated rats: a model for Smith-Lemli-Opitz syndrome. *Journal of lipid*
898 *research*, 52(10), 1810-1820.

899 Xu, L., Sheflin, L.G., Porter, N.A., & Fliesler, S.J. (2012). 7-Dehydrocholesterol-derived oxysterols and
 900 retinal degeneration in a rat model of Smith–Lemli–Opitz syndrome. *Biochimica et Biophysica Acta*
 901 *(BBA)-Molecular and Cell Biology of Lipids*, 1821(6), 877-883.
 902 Yang T., Espenshade P. J., Wright M. E., Yabe D., Gong Y., Aebersold R., Goldstein J. L., Brown M. S.
 903 (2002) Crucial step in cholesterol homeostasis. Sterols promote binding of SCAP to INSIG-1, a
 904 membrane protein that facilitates retention of SREBPs in ER. *Cell* 110, 489–500.
 905 Yang, C., Xie, L., Gu, Q., Qiu, Q., Wu, X., & Yin, L. (2019). 7-Ketocholesterol disturbs RPE cells
 906 phagocytosis of the outer segment of photoreceptor and induces inflammation through ERK
 907 signaling pathway. *Experimental eye research*, 189, 107849.
 908 Zhao, Y., Mazzone, T. (1999). LDL receptor binds newly synthesized apoE in macrophages. A
 909 precursor pool for apoe secretion. *Journal of Lipid Research*, 40, 1029–1035.
 910 Zheng, W., Reem, R.E., Omarova, S., Huang, S., DiPatre, P.L., Charvet, C.D., ... Pikuleva, I.A. (2012).
 911 Spatial distribution of the pathways of cholesterol homeostasis in human retina. *PloS one*, 7(5).
 912
 913
 914
 915
 916
 917
 918
 919
 920
 921
 922
 923
 924

Figure legends

Figure 1 Structure of oxysterols generated in retina and RPE. The main oxysterols that are oxidized from cholesterol or its precursors in either enzymatic or non-enzymatic way. The enzymatic oxygenation reactions are mainly mediated by cytochrome P-450 family (green arrows). The non-enzymatic oxysterol-generating reactions are catalysed by reactive oxygen species (ROS) (red arrow).


Figure 2 Retinal structure and localization of oxysterols and major oxysterol-metabolism enzymes. (A) Structure of human eye and retina. Left panel: Human retina (yellow) is located on the rear hemisphere of the eyeball. From outermost to innermost the layers are the sclera, the choroid and the retina. The macula is an oval pigmented area located at the central retina where the highest concentration of cones is found. Retinal blood vessels enter the retina in conjunction with the optic nerve, with branches distributing in the choroid layer. The cornea is a transparent layer at the front of the eye, lying anterior to the iris and pupil. Right panel: human retina consists of different types of cells: retinal pigment epithelium (RPE) cells, rod photoreceptors, cone photoreceptors, bipolar cells, horizontal cells, amacrine cells, ganglion cells and Müller cells. They form several layers of the retina (from outermost to innermost): outer nuclear layer (ONL), outer plexiform layer (OPL), inner nuclear layer (INL), inner plexiform layer (IPL), ganglion cell layer (GCL) and nerve fibre layer (NFL). Müller cells span the entire retina. (B) Distribution of oxysterol-producing enzymes and presentation of main oxysterols in the retina.

Figure 3 TSPO-mediated cholesterol efflux in retinal pigment epithelial (RPE) cells. The RPE uptakes low-density lipoproteins (LDL) from the choroidal circulation mainly via LDL receptor (LDLR). Cholesterol is released from LDL degradation in the RPE and delivered to the mitochondria, where cholesterol is transported from mitochondrial outer membrane (MOM) to mitochondrial inner membrane (MIM) by TSPO protein complex containing TSPO, steroidogenic acute regulatory protein (StAR, not shown), the voltage dependent anion channel (VDAC), the adenine nucleotide


translocator (ANT) and other regulatory proteins (not shown). Oxysterol is generated by CYP27A1 to 27-hydroxycholesterol (27-OHC), which upregulates expression of cholesterol transports (e.g. ABCA1 and ABCG1A) via activation of the liver-X receptor (LXR) signalling pathway and enhances cholesterol efflux. Ligands can bind TSPO, enhance mitochondrial cholesterol trafficking and promote cholesterol efflux. BrM, Bruch's membrane; MOM: Mitochondrial outer membrane; MIM: Mitochondrial inner membrane; MM: Mitochondrial matrix; IPM, interphotoreceptor matrix; ApoA1: Apolipoprotein A1; ApoE: Apolipoprotein E; ABCA1: ATP-binding cassette transporter member 1; ABCG1: ATP-binding cassette sub-family G member 1.

Figure1


Figure 1



A



B



997

998

999

1000


1001

1002

1003

1004

1005 **Figure 3**



1016 Table S1 Abbreviation and nomenclature of oxysterols

Abbreviation	Common name	Systemic name	Lipid Map ID
4 β -HC	4 β -hydroxycholesterol	Cholest-5-en-3 β ,4 β -diol	LMST01010014
7 α -HC	7 α -hydroxycholesterol	Cholest-5-en-3 β ,7 α -diol	LMST01010013
7-DHC	7-dehydrocholesterol	Cholesta-5,7-dien-3 β -ol	LMST01010069
7-KC	7-ketocholesterol	7-oxo-Cholest-5-en-3 β -ol	LMST01010049
7-KCh-27OH	3 β ,27-dihydroxy-5-cholesten-7-one	(25R)-7-oxo-cholest-5-en-3 β ,27-diol	LMST04030180
7 α ,24S-diHC	7 α ,24(S)-dihydroxycholesterol	Cholest-5-en-3 β ,7 α ,24-triol	LMST04030174
7 α ,25-diHC	7 α ,25-dihydroxycholesterol	Cholest-5-en-3 β ,7 α ,25-triol	LMST04030166
7 α ,27-diHC	7 α ,27-dihydroxycholesterol	5-Cholestene-3 β ,7 α ,26-triol	LMST04030081
20R,20R-diHC	(20R,22R)-20,22-dihydroxycholesterol	(3 β)-cholest-5-ene-3,20,22-triol	LMST01010200
22R-HC	22(R)-Hydroxycholesterol	Cholest-5-en-3 β ,22R-diol	LMST01010086
24S-HC	24(S)-Hydroxycholesterol	Cholest-5-en-3 β ,24S-diol	LMST01010019
24S,25-EC	24(S),25-Epoxycholesterol	24S,25-Epoxy-cholest-5-en-3 β -ol	LMST01010012
25-HC	25-Hydroxycholesterol	Cholest-5-en-3 β ,25-diol	LMST01010018

1017
1018
1019
1020